ITAMIN C SUPPLEMENT

Composition

Each ml contains: vitamin C (ascorbic acid) 200 mg.

Indications

- Prevention or treatment of vitamin C deficiencies in farm animals, including increased disease incidence.
- Convalescence
- Improve the immune response.
- Prevention against copper-induced damage in erythrocyte and liver of broiler chickens.
- Prevention against infectious bursal disease, malabsorption syndrome, adenovirus infections, chicken anaemia agent, mycotoxins or coccidiosis.
- Prevention and treatment of stress caused by vaccination, diseases, transport, high humidity, high temperatures or extreme temperature changes.


Batch No and Expiry date: See package.

Manufactured by:



SUPER'S DIANA, S.L. Ctra C-17 Km 17 08150 Parets del Valles SUPER'S DIANA (Barcelona) SPAIN. www.supersdiana.com

(FOR VETERINARY USE ONLY)



Cattle, sheep, goats and poultry.

Dosage and route of administration

Oral use in drinking water.

- Poultry: 1 ml/10 litres drinking water for 3 to 5 days.
- Calves and cows, goats and sheep: 1 ml/50 kg bw for 3 to 5 days.

Withdrawal period

Zero days.

Storage conditions

Store below 30 °C. Protect from light.

Manufacturer's inscription number: αESP08100107



Exclusive Distributor in Pakistan:



Opp. Kings Mall, Near Wapda Town, G.T. Road, Gujranwala-Pakistan. Ph: +92 55-4284177, 42824188

